

**Titre:** The addition of the collagen binding domain of fibronectin potentiates the biochemical availability of basic fibroblast growth factor for the culture of induced pluripotent stem cells  
**Title:**

**Auteurs:** Cyril Addi, Frédéric Murschel, Benoit Liberelle, Nesrine Riahi, Olivier Feraud, & Gregory De Crescenzo  
**Authors:**

**Date:** 2016

**Type:** Communication de conférence / Conference or Workshop Item

**Référence:** Addi, C., Murschel, F., Liberelle, B., Riahi, N., Feraud, O., & De Crescenzo, G. (2016, May). The addition of the collagen binding domain of fibronectin potentiates the biochemical availability of basic fibroblast growth factor for the culture of induced pluripotent stem cells [Poster]. 10th World Biomaterials Congress, Montréal, Québec. Published in Frontiers in Bioengineering and Biotechnology. <https://doi.org/10.3389/conf.fbioe.2016.01.00338>  
**Citation:**

## Document en libre accès dans PolyPublie

Open Access document in PolyPublie

**URL de PolyPublie:** <https://publications.polymtl.ca/4835/>  
**PolyPublie URL:**

**Version:** Version officielle de l'éditeur / Published version  
Révisé par les pairs / Refereed

**Conditions d'utilisation:** CC BY  
**Terms of Use:**

## Document publié chez l'éditeur officiel

Document issued by the official publisher

**Nom de la conférence:** 10th World Biomaterials Congress  
**Conference Name:**

**Date et lieu:** 2016-05-17 - 2016-05-22, Montréal, Québec  
**Date and Location:**

**Maison d'édition:** Frontiers Media S.A.  
**Publisher:**

**URL officiel:** <https://doi.org/10.3389/conf.fbioe.2016.01.00338>  
**Official URL:**

**Mention légale:**  
**Legal notice:**


## EVENT ABSTRACT

[« Back to Event](#)

# The addition of the collagen binding domain of fibronectin potentiates the biochemical availability of basic fibroblast growth factor for the culture of induced pluripotent stem cells

Cyril Addi<sup>1, 2</sup>, Frederic Murschel<sup>1</sup>, Benoit Liberelle<sup>1</sup>, Nesrine Riahi<sup>1</sup>, Olivier Feraud<sup>2</sup> and Gregory De Crescenzo<sup>1</sup><sup>1</sup> Ecole Polytechnique de Montreal, Department of Chemical Engineering, Canada<sup>2</sup> Université Paris-Sud, Hôpital Paul Brousse, INSERM U935, ESteam Paris-Sud, France


**Introduction:** The culture of various cell types including induced pluripotent stem cells (iPSC), requires the use of growth factors such as bFGF (basic Fibroblast Growth Factor)<sup>[1],[2]</sup>. These factors are generally added to the medium in their diffusive form and have a very short half-life<sup>[3]</sup>: they disappear quickly from the culture medium and must be replaced at high cost. We here explore the stable and oriented tethering of bFGF on a cell culture substrate, namely gelatin (denatured collagen). Our strategy relies on the expression of two fusion proteins being labelled with two distinct peptides (the E and Kcoils) that bind to each other with great affinity and specificity<sup>[4]</sup>. That is, a chimera corresponding to bFGF fused to the Ecoil peptide on the one hand, and the collagen-binding domain (CBD) of fibronectin fused to the Kcoil peptide on the other hand (**Fig. 1**).



**Figure 1:** Schematic illustration of Ecoil-bFGF tethered in an oriented manner on a gelatin-coated surface functionalized with CBD-Kcoil.

**Materials and Methods:** The Ecoil-bFGF fusion protein was produced in *E. coli* and purified by Ni-NTA affinity chromatography followed by TEV protease cleavage and ion exchange chromatography. The CBD-Kcoil fusion protein was produced in *E. coli* and purified by solubilizing inclusions bodies in an 8 M urea buffer, dialyzing against PBS, and then followed by a Ni-NTA affinity chromatography. The ability of Ecoil-bFGF protein to bind to Kcoil was assessed by surface plasmon resonance (SPR), using a Kcoil-coated chip while its ability to bind to its cognate receptor was tested by subsequent injections of a recombinant chimera of the FGF Receptor (FGFR1-Fc). Finally, the suitability of the proposed approach for cell culture will be soon assessed by cultivating iPSCs on a gelatin-coated substrate to be functionalized with CBD-Kcoil/Ecoil-bFGF.

**Results and Discussion:** Both fusion proteins were produced and purified with high purity. The yield of the purification of Ecoil-bFGF was assessed by ELISA and reached 1.1 mg per liter of bacteria. On the one hand, SPR analyses (**Fig. 2**) demonstrated that Ecoil-bFGF specifically bound to a Kcoil-coated chip, and subsequently interacted with FGFR1-Fc with high affinity ( $K_D = 1.8$  nM), in good agreement with the literature<sup>[5]</sup>. On the other hand, the bioactivity of CBD-Kcoil was assessed via an Enzyme-Linked Immunosorbent Assay (ELISA) against bFGF: gelatin-coated substrates were incubated with CBD-Kcoil then Ecoil-bFGF, and the formation of the gelatin/CBD-Kcoil/Ecoil-bFGF complex was demonstrated by the specific recruitment of anti-bFGF antibodies. Altogether, the data indicated that our novel strategy enabled the specific capture of bioactive bFGF via collagen/CBD and coiled-coil interactions.



**Figure 2:** SPR biosensor-based assays demonstrating the bioavailability of Ecoil and bFGF moieties. Control-corrected sensorgrams are presented. (A) Injection of 10 nM Ecoil-bFGF over Kcoil. (B) Injection of increasing concentrations of FGFR1-Fc over Ecoil-bFGF immobilized by coiled-coil interactions.

**Conclusion:** Providing iPSCs with bioactive bFGF in a non-diffusible form may significantly decrease expenses related to cell culture. Furthermore, the modular approach we undertook may make this strategy amenable to the development of more complex scaffolds combining several growth factors (bFGF, EGF, VEGF ...), and thus pave the way to the development of novel approaches in the field of tissue engineering.

*This project was supported by the Natural Sciences and Engineering Research Council of Canada (NSERC), CREATE program - MEDITIS scholarship.*

## References:

- [1] M. Amit, M. K. Carpenter, M. S. Inokuma, C.-P. Chiu, C. P. Harris, M. A. Waknitz, J. Itskovitz-Eldor, and J. A. Thomson, “Clonally Derived Human Embryonic Stem Cell Lines Maintain Pluripotency and Proliferative Potential for Prolonged Periods of Culture,” *Dev. Biol.*, vol. 227, no. 2, pp. 271–278, Nov. 2000.
- [2] K. Takahashi, K. Tanabe, M. Ohnuki, M. Narita, T. Ichisaka, K. Tomoda, and S. Yamanaka, “Induction of Pluripotent Stem Cells from Adult Human Fibroblasts by Defined Factors,” *Cell*, vol. 131, no. 5, pp. 861–872, Nov. 2007.
- [3] G. Chen, D. R. Gulbranson, P. Yu, Z. Hou, and J. A. Thomson, “Thermal Stability of Fibroblast Growth Factor Protein Is a Determinant Factor in Regulating Self-Renewal, Differentiation, and Reprogramming in Human Pluripotent Stem Cells,” *Stem Cells Dayt. Ohio*, vol. 30, no. 4, pp. 623–630, Apr. 2012.
- [4] F. Murschel, B. Liberelle, G. St-Laurent, M. Jolicoeur, Y. Durocher, and G. De Crescenzo, “Coiled-coil-mediated grafting of bioactive vascular endothelial growth factor,” *Acta Biomater.*, vol. 9, no. 6, pp. 6806–6813, Jun. 2013.
- [5] X. Lin, K. Takahashi, S. L. Campion, Y. Liu, G. G. Gustavsen, L. A. Peña, and P. O. Zamora, “Synthetic peptide F2A4-K-NS mimics fibroblast growth factor-2 in vitro and is angiogenic in vivo,” *Int. J. Mol. Med.*, vol. 17, no. 5, pp. 833–839, May 2006.

**Keywords:** stem cell, biosensing, growth factor, biofunctionalization **Conference:** 10th World Biomaterials Congress, Montréal, Canada, 17 May - 22 May, 2016.

**Presentation Type:** Poster **Topic:** Protein interactions with biomaterials

**Citation:** Addi C, Murschel F, Liberelle B, Riahi N, Feraud O and De Crescenzo G (2016). The addition of the collagen binding domain of fibronectin potentiates the biochemical availability of basic fibroblast growth factor for the culture of induced pluripotent stem cells. *Front. Bioeng. Biotechnol. Conference Abstract: 10th World Biomaterials Congress*. doi: 10.3389/conf.FBIOE.2016.01.00338

**Copyright:** The abstracts in this collection have not been subject to any Frontiers peer review or checks, and are not endorsed by Frontiers. They are made available through the Frontiers publishing platform as a service to conference organizers and presenters.

The copyright in the individual abstracts is owned by the author of each abstract or his/her employer unless otherwise stated.

Each abstract, as well as the collection of abstracts, are published under a Creative Commons CC-BY 4.0 (attribution) licence (<https://creativecommons.org/licenses/by/4.0/>) and may thus be reproduced, translated, adapted and be the subject of derivative works provided the authors and Frontiers are attributed.

For Frontiers' terms and conditions please see <https://www.frontiersin.org/legal/terms-and-conditions>. Received: 27 Mar 2016; Published Online: 30 Mar 2016.